

# Seminar: DNA Hybridization Detection

Chris Harris

Chemistry Department

University of Maine

March 21, 2006

# Outline

Ruthenium project

General principles

Literature review



Experimental data

Future prospects

Conclusion





# Role of messenger RNA in biological systems [5]




Transcription versus reverse transcription


# Discovery of RNA interference (RNAi) [4]



Messenger RNA for purple color inserted in petunia seeds



# Promise of RNA interference (RNAi)






Revolutionary drug therapy allows one to shut off genes in hereditary diseases:



Diabetes  
Alzheimers  
Breast Cancer  
Depression



# ElectroChemiluminescence of $\text{Ru}(\text{bpy})_3^{+2}$ [1]




# Catalytic effects on cyclic voltammograms of $\text{Ru}(\text{bpy})_3^{+2}$ [1]



Glassy Carbon versus Platinum material


## How DNA labeling works [2]



### General methodology:


- 1) Attach probe ssDNA to magnetic bead
- 2) Attach target ssDNA to polystyrene bead
- 3) Hybridize DNA strands in aqueous media
- 4) Isolate hybridized pairs with magnet
- 5) Perform ECL in acetonitrile

## Fluorescence of polystyrene beads [2]




(a)  $\text{Ru}(\text{bpy})_3^{2+}$ ; (b)  $\text{Ru}(\text{bpy})_3^{2+} + \text{avidin}$

## Scanning electron image [2]




Polystyrene balls surrounded by magnetic beads after hybridization

## Electrochemical behavior [2]




(a) CV of  $\text{Ru}(\text{bpy})_3^{2+}$  + acetyl nitrile + salt + tripropyl amine; (b) ECL response; (c) CV in the absence of tripropyl amine [I x 10]

## ECL intensity as a function of tripropyl amine concentration [2]




$\text{Ru}(\text{bpy})_3^{2+}$ , acetylnitrile, and salt present

# ECL intensity as a function of trifluoroacetic acid addition: pH optimization [2]




$\text{Ru}(\text{bpy})_3^{2+}$ , acetylitrile, salt, and tripropyl  
amine present

## Eradication of tripropyl amine ECL [2]




- (a) Tripropyl amine + salt + acetonitrile;
- (b) Add trifluoroacetic acid; (c) Add 1 vol% water; (d) Salt + acetonitrile alone

## ECL intensity in relation to $\text{Ru}(\text{bpy})_3^{2+}$ concentration [2]





Tripropyl amine, trifluoroacetic acid, salt,  
acetonitrile, and water present

## ECL intensity as a function of polystyrene ball quantity [2]



$\text{Ru(bpy)}_3^{2+}$  labeling

## ECL detection of DNA hybridization [2]





- (a) Small magnetic beads and high population ratio;  
(b) More favorable magnetic bead size and population ratio

## Advantages over competing ECL techniques: [2]

- Dual phases: aqueous DNA coupling in combination with acetonitrile analysis.
- High selectivity/ low detection limit: starting as low as 1.0 fM concentration for target DNA.
- Unprecedented stability: once  $\text{Ru}(\text{bpy})_3^{2+}$  label is released into acetonitrile, ECL measurement can be performed repeatedly.


# Chemical processes involved in typical ECL systems [3]



where: TPA = tripropyl amine

DPA = dipropyl amine

# Cyclic voltammogram of $\text{Ru}(\text{bpy})_3^{2+}$




2<sup>nd</sup> Cycle

Scanrate= 200 mV/s

Interdigitated glassy carbon electrode

# Cyclic voltammogram of $\text{Ru}(\text{bpy})_3^{2+}$



## Research Frontiers:

- Carbon rod or Pt electrode in acetonitrile using pulses.
- Rotating electrode in acetonitrile to find lifetime of  $\text{Ru}(\text{bpy})_3^{+1}$ .
- Try vertically patterned electrodes rather than interdigitated configuration.

# Vertically integrated electrodes

1. Deposit PECVD  $\text{Si}_3\text{N}_4$  on Carbon Coated Wafers

Note: Silicon wafer are coated with oxide/nitride before Carbon deposition



2. Spin/Pattern Photoresist




3. Evaporate Platinum



4. Strip Resist



5. Etch  $\text{Si}_3\text{N}_4$  using a  $\text{SF}_6$  plasma



Deposition steps and top view of vertical matrix

## Conclusions:

- Whole new drug development scheme on the horizon which identifies genes attributed to disease, then shuts off expression with messenger RNA.
- Relative to its rival fluorescence, ECL eliminates the excitation light required, enhancing sensitivity and reducing cost.
- With the base pair selectivity ECL offers, DNA matching could evolve into sequencing.

## References

- [1] G. C. Fiaccabruno, et al., *Analytical Chemistry*, **70**, 4157 (1998).
- [2] W. Miao and A. J. Bard, *Analytical Chemistry*, **76**, 5379 (2004).
- [3] J. K. Leland and M. J. Powell, *Journal of Electrochemistry Society*, **137**, 3127 (1990).
- [4] C. Napoli, C. Lemieux, and R. Jorgensen, *Plant Cell*, **2**, 279 (1990).
- [5] <http://en.wikipedia.org>